

There are two main types of mountain biking injuries, those caused by falling off – acute or traumatic injuries which are discussed in another leaflet, and those issues caused by overtraining, biomechanical stresses, often due to muscle imbalances, and incorrect bike set-up. It is particularly important for the sake of future injury prevention to identify the root cause and address this ASAP, otherwise the problem will persist. It's definitely worth seeking the advice of a physical therapist as well as someone trained specifically in bike set up.

1 KNEE PAIN

One of the most common riding knee complaints is pain in the kneecap. This is most likely to be patellofemoral pain syndrome (PFPS). PFPS is often worse when walking up and down hills/stairs or sitting for long periods of time. It may include wasting of the quadriceps (thigh) muscles if the injury is an old one and tight muscles around the knee joint.

PFPS occurs when the patella (kneecap) rubs on the femur (thigh) bone underneath. It is believed that incorrect tracking (gliding) of the patella over the femur is a significant factor and results in damage to the cartilage underneath the patella. The cause may be from external factors like an increase in training, the seat being too low or riding for too long in big gears. Internal factors such as poor patella tracking may result from excessive pronation (flat foot), rotation of the lower leg and tight or weak muscles around the thigh and pelvis.

TIP: Try variation in your pedal cadence – breaking up your training by pedalling in a high cadence (90 – 120 revs per minute) can help prevent injury. Raising the seat if it's too low will reduce the amount of knee flexion and utilise more of the hamstring and gluteal (buttock) muscles thereby off-loading the quadriceps muscle and patellar tendon. If your foot rocks from side to side, use an insert to stabilise the rear of your foot, thereby reducing strain on the knee and increasing efficiency.

2 BACK PAIN

After knees, the back is probably one of the biggest causes of pain for cyclists, with lack of flexibility and bad posture generally the cause. Hunching forward on your bike, and probably also at work, places strain on your spine, loading structures

OVERUSE INJURIES IN MOUNTAIN BIKING AND HOW TO AVOID THEM

for prolonged periods of time. Cyclists' back pain is often due to mechanical factors. Have your bike properly fitted to your body, then look at your body. Lack of flexibility, such as excessive hamstring and hip flexor tightness can contribute to low back pain. Differences in leg length are common mechanical problems leading to imbalances in the spine. Core strength is very important to avoid low back pain. Core strength comes from a collection of deep muscles both big and small that work together to give you core lumbar and pelvic stability.

TIP: Low back pain may arise in cyclists that push big gears, especially while climbing. The angle of your back in relation to the bike can increase or decrease the strain on your back. Consider alternating climbing positions by standing up, changing the angle of your back, especially during long rides or climbs or talk to your physical therapist about a back rehabilitation

programme to focus on strengthening your core muscles.

3 NECK PAIN

Neck pain from cycling usually stems from poor posture and weak muscles. Pain caused by neck hyperextension is made worse by positional issues on the bike, combined with lack of flexibility. Just as you have core stabilisers around your lower back, you have stabiliser muscles called deep neck flexors around your neck to hold your head up. When your neck stabilisers are weak or fatigue quickly it is left to the trapezius muscle (that goes from the base of your skull to the tip of the shoulder) to support your head as you lean forward. And when these 'stand-in' muscles fatigue you can experience pain in the back and sides of your neck. Restore balance by keeping the neck muscles loose and relaxed through a routine of strengthening and stretching exercises.

TIP: Change your posture on the bike. If you're reaching too far forward, or your handlebars are too low, shorten the stem to shorten your reach. Raise your bars and riding more upright will reduce the strain you're putting on your back and neck. Don't forget to change your hand positions at regular intervals, and sit up on the bike to stretch, straightening out your neck and back to vary the loads on the different muscle groups.

4 ILIOTIBIAL BAND (ITB) PAIN

While it is more commonly known as "runner's knee," ITB syndrome is another common cycling injury. ITB pain is typically associated with prolonged, repetitive activity. Symptoms include pain on the outside of the knee, tenderness and sometimes swelling. In some cases, pain is felt simply walking or going up and down stairs. You may feel stiff or tight after periods of inactivity.

The ITB is a tendinous fascial band that originates on the iliac crest (hipbone) and attaches just below on the outside of the knee. As your knee bends and straightens repeatedly, the band can become inflamed from rubbing over bony condyles. Other contributing factors may include tightness of thigh, hip and buttock muscles as well as weak pelvic stabilising muscles.

TIP: Ensure your seat is at the optimal height for your body. If you pronate (are flat footed) you may need orthotics or a wedge inside your cycling shoe to stop the leg and knee rotating inwards putting further tension on the ITB. Tucking the knees in too tight, to increase streamline, will also add tension and greater friction to the ITB.

5 ACHILLES TENDON PAIN

The Achilles tendon is the tendon at the back of the ankle, connecting the gastrocnemius (calf) muscle to the heel. If your Achilles is sore during or after riding you may have Achilles tendinopathy. Inflammation, micro-tears or compromised blood flow, often caused by overuse, could put a stop to your riding season. There is a whole host of stretching and strengthening

options available from your therapist.

TIP: Having your saddle too high keeps the foot plantarflexed (toe pointed down), causing constant contraction of the calf muscles and load on the tendon. Lowering your seat and making sure that your cleats aren't pushed all the way forward towards the toe will help to even out what muscles you're using to pedal.

6 HIP PAIN

Possible causes of hip pain in cyclists include bursitis, snapping hip syndrome, impingement syndrome, labral tears or piriformis syndrome. Although the diagnoses may vary, the causes of cycling hip injuries are usually similar and involve over-training, pushing excessively high gears and muscle imbalances. For example, piriformis syndrome is caused by overuse of the gluteal (buttock) muscles, which results in a weak, tight piriformis muscle that can cause sciatica.

TIP: Address the underlying muscle imbalances - by strengthening a muscle, the tightness will ease off and often the pain will disappear too. Gear back and increase your cadence to take pressure off your hips.

7 BURNING FEET

Painful burning of the ball of the foot (a.k.a. "hot foot," or metatarsalgia) is usually a result of hot weather and/ or poorly fitting shoes on long, hilly rides. Pressure can pinch nerves in one or both feet.

TIP: Make sure your cycling shoes are wide enough and not 'squishing' your feet. Move your cleats a few millimetres closer to the heel of the shoe to take pressure off your forefoot. Or switch to a larger-platform pedal to more evenly distribute the pressure across your feet. Cushion the blow by adding some supportive insoles and choose thermo-regulating socks.

8 HAND PAIN

'Handlebar Palsy' is a name given to a condition suffered by cyclists caused by compression of the ulnar nerve at the wrist against the handlebar. It often comes on after long rides, and is not just due to the pressure from your weight but also the transmission of road 'buzz' and vibration through the bars. Symptoms include numbness, tingling and weakness over the outside of the hand, little finger and outer half of the ring finger. A feeling of clumsiness in the hand is often reported and pain may be present when moving the wrist.

TIP: Treat the cause of the problem. In cyclists, this may mean checking the bike set up, often you need to shorten your reach, that way more of your bodyweight is borne by the saddle than your hands. Gel padded gloves and padded bars can give cushioning and reduce the pressure on the nerve.

CONCLUSION

Many cycling overuse injuries can be prevented. Be assessed for underlying ALIGNMENT issues such as muscle weaknesses, flexibility, leg length discrepancies and being flat-footed. Then check you have the correct BIKE set up, correct technique and cycling posture. Finally, CONDITION yourself with appropriate strengthening and stretching exercises and progress your training moderately and always, always listen to your body.



The information contained in this article is intended as general guidance and information only and should not be relied upon as a basis for planning individual medical care or as a substitute for specialist medical advice in each individual case. ©Co-Kinetic 2020